arXiv:1408.0569v1 [math.PR] 4 Aug 2014

# Wong-Zakai Approximations of Backward Doubly Stochastic Doubly Backward Differential Equations

Ying Hu[1], Anis Matoussi[2] and Tusheng Zhang[3]

August 5, 2014

**Abstract**

In this paper we obtain a Wong-Zakai approximation to solutions of backward doubly stochastic differential equations.

**AMS Subject Classification:** Primary 60H15 Secondary 93E20, 35R60.

## 1 Framework and Introduction

Let $\{W_t, 0 \leq t \leq T\}$ and $\{B_t, 0 \leq t \leq T\}$ be two independent standard Brownian motions on a probability space $(\Omega, \mathcal{F}, P)$. Let $\mathcal{N}$ denote the class of $P$-null sets. For each $t \in [0,T]$, we define

$$\mathcal{F}_t = \mathcal{F}_t^W \vee \mathcal{F}_{t,T}^B,$$

where for any process $\{\eta_t\}$, $\mathcal{F}_{s,t}^\eta = \sigma\{\eta_r - \eta_s; s \leq r \leq t\} \vee \mathcal{N}$,$\mathcal{F}_t^\eta = \mathcal{F}_{0,t}^\eta$. Let $f : R \times R \to R$ be a bounded measurable function satisfying

**(H.1)**

$$|f(y_1, z_1) - f(y_2, z_2)| \leq c(|y_1 - y_2| + |z_1 - z_2|)$$

Let $g \in C_b^1(R)$. For $n \geq 1$, define the linear interpolation $B^n$ of $B$ as

$$B_t^n = B_{\frac{k+2}{2^n}} + 2^n(t - \frac{k+1}{2^n})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}}), \quad \text{for} \quad t \in [\frac{k}{2^n}, \frac{k+1}{2^n}]. \tag{1.1}$$

Let $\xi \in L^2(\Omega)$ be $\mathcal{F}_T$-measurable. Consider the following backward doubly stochastic differential equations(BDSDE):

$$\begin{aligned} Y_t &= \xi + \int_t^T f(Y_s, Z_s)ds + \int_t^T g(Y_s)dB_s \\ &\quad + \frac{1}{2}\int_t^T gg'(Y_s)ds - \int_t^T Z_s dW_s. \end{aligned} \tag{1.2}$$

[1] Université de Rennes 1, campus Beaulieu, 35042 Rennes Cedex, France

[2] Laboratoire Manceau de Mathématiques, Université du Maine, Avenue Olivier Messiaen, 72 085 LE MANS Cedex, France

[3] School of Mathematics, University of Manchester, Oxford Road, Manchester M13 9PL, England, U.K. Email: tusheng.zhang@manchester.ac.uk

$$Y_t^n = \xi + \int_t^T f(Y_s^n, Z_s^n)ds + \int_t^T g(Y_s^n)dB_s^n - \int_t^T Z_s^n dW_s. \tag{1.3}$$

Here $dB_s$,$dB_s^n$ stand for the backward integrals. In the sequel, we will use $\overrightarrow{ds}$ to indicate the backward integral against the Lebesgue measure.

Backward doubly stochastic differential equations was first studied by Pardoux and Peng in [PP]. It is now a powerful tool to study stochastic partial differential equations with singular coefficients. Our purpose of this paper is to obtain the convergence of the Wong-Zakai approximation to the backward doubly stochastic differential equations, namely we will prove that $(Y^n, Z^n)$ converges to $(Y, Z)$ in $L^2$. The convergence of Wong-Zakai approximations to stochastic differential equations is now well known, see e.g. [IW]. Because of the nature of the BDSDEs, the integrand $Z^n, Z$ in the stochastic integral against Brownian motion are also part of the solutions. This makes the problem drastically different from the Wong-Zakai approximation for stochastic different equations. Another difficulty comes from the fact that the Hölder type estimate

$$E[|Y_t^n - Y_s^n|^p] \leq C|t-s|^\alpha$$

is no longer available. We overcome this by carefully exploiting the independence of the two Brownian motions $B$ and $W$.

The application of our results to stochastic partial differential equations will be discussed in a forthcoming paper.

# 2 Main results

The following is an priori estimate for the family $\{(Y^n, Z^n), n \geq 1\}$.

**Proposition 2.1** *There exists a constant $C$ such that*

$$\sup_n \sup_{0\leq t\leq T} \{E[(Y_t^n)^2] + E[\int_t^T (Z_s^n)^2 ds]\} \leq C. \tag{2.4}$$

**Proof**. By Ito's formula, we have

$$\begin{aligned} & (Y_t^n)^2 + \int_t^T (Z_s^n)^2 ds \\ = \; & (\xi)^2 + 2\int_t^T Y_s^n f(Y_s^n, Z_s^n)ds + 2\int_t^T Y_s^n g(Y_s^n)dB_s^n \\ & -2\int_t^T Y_s^n Z_s^n dW_s. \end{aligned} \tag{2.5}$$

For $s \in [\frac{k}{2^n}, \frac{k+1}{2^n}]$, set $s^+ = \frac{k+2}{2^n}$ and $s^- = \frac{k-1}{2^n}$. In view of (H.1), it is easy to see that there exists $C_1 > 0$ such that

$$2\int_t^T Y_s^n f(Y_s^n, Z_s^n)ds \le C_1 \int_t^T (Y_s^n)^2 ds + \frac{1}{4}\int_t^T (Z_s^n)^2 ds + C_1. \tag{2.6}$$

Now, the third term on the right side of (2.5) can be written as

$$\begin{aligned} 2\int_t^T Y_s^n g(Y_s^n)dB_s^n &= 2\int_t^T Y_{s^+}^n g(Y_{s^+}^n)dB_s^n \\ &+ 2\int_t^T (Y_s^n - Y_{s^+}^n)g(Y_s^n)dB_s^n \\ &+ 2\int_t^T Y_{s^+}^n (g(Y_s^n) - g(Y_{s^+}^n))dB_s^n \\ &:= I_1 + I_2 + I_3. \end{aligned} \tag{2.7}$$

As a stochastic integral, we have $E[I_1] = 0$. By the equation (1.3) it follows that

$$\begin{aligned} I_2 &= 2\int_t^T (\int_s^{s^+} f(Y_u^n, Z_u^n)du)g(Y_s^n)dB_s^n \\ &+ 2\int_t^T (\int_s^{s^+} g(Y_u^n)dB_u^n)g(Y_s^n)dB_s^n \\ &- 2\int_t^T (\int_s^{s^+} Z_u^n dW_u)g(Y_s^n)dB_s^n \\ &:= I_{2.1} + I_{2.2} + I_{2.3}. \end{aligned} \tag{2.8}$$

By the boundedness of $f$ and $g$, we have

$$\begin{aligned} E[I_{2.1}] &\le C\int_t^T (\int_s^{s^+} du)E[|\dot{B}_s^n|]ds \\ &\le C(\frac{1}{2^n})^{\frac{1}{2}}, \end{aligned} \tag{2.9}$$

and

$$\begin{aligned} E[I_{2.2}] &\le CE[\int_t^T (\int_s^{s^+} |\dot{B}_u^n|du)|\dot{B}_s^n|ds] \\ &\le C\int_t^T ds\int_s^{s^+} duE[|\dot{B}_u^n||\dot{B}_s^n|] \\ &\le C\int_t^T ds\int_s^{s^+} du(E[|\dot{B}_u^n|^2])^{\frac{1}{2}}(E[|\dot{B}_s^n|^2])^{\frac{1}{2}} \\ &\le C\int_t^T ds\int_s^{s^+} du(2^n)^{\frac{1}{2}}(2^n)^{\frac{1}{2}} \le C, \end{aligned} \tag{2.10}$$

where $C$ is a constant independent of $n$. For the term $I_{2.3}$, we have

$$\begin{aligned} E[I_{2.3}] &= 2\int_t^T E[(\int_s^{s^+} Z_u^n dW_u) g(Y_s^n)\dot{B}_s^n]ds \\ &= 2\int_t^T E[g(Y_s^n)\dot{B}_s^n E[(\int_s^{s^+} Z_u^n dW_u)|\mathcal{F}_s]]ds \\ &= 0. \end{aligned} \tag{2.11}$$

Putting together (2.9),(2.10),(2.11) we get

$$\sup_n E[I_2] \leq C, \tag{2.12}$$

for some constant $C$. To bound $I_3$ in (2.7), we write

$$\begin{aligned} I_3 &= 2\int_t^T Y_{s^+}^n \int_0^1 d\lambda g'(Y_{s^+}^n + \lambda(Y_s^n - Y_{s^+}^n))(Y_s^n - Y_{s^+}^n)dB_s^n \\ &= 2\int_t^T Y_{s^+}^n \int_0^1 d\lambda g'(Y_{s^+}^n + \lambda(Y_s^n - Y_{s^+}^n))(\int_s^{s^+} f(Y_u^n, Z_u^n)du)dB_s^n \\ &+ 2\int_t^T Y_{s^+}^n \int_0^1 d\lambda g'(Y_{s^+}^n + \lambda(Y_s^n - Y_{s^+}^n))(\int_s^{s^+} g(Y_u^n)dB_u^n)dB_s^n \\ &- 2\int_t^T Y_{s^+}^n \int_0^1 d\lambda g'(Y_{s^+}^n + \lambda(Y_s^n - Y_{s^+}^n))(\int_s^{s^+} Z_u^n dW_u)dB_s^n \\ &:= I_{3.1} + I_{3.2} + I_{3.3}. \end{aligned} \tag{2.13}$$

For $I_{3.1}$, we have

$$\begin{aligned} E[I_{3.1}] &\leq C\int_t^T E[|Y_{s^+}^n|\frac{1}{2^n}|\dot{B}_s^n|]ds \\ &\leq C\int_t^T E[(Y_{s^+}^n)^2]ds + C\int_t^T E[(\frac{1}{2^n})^2|\dot{B}_s^n|^2]ds \\ &\leq C\int_t^T E[(Y_{s^+}^n)^2]ds + C. \end{aligned} \tag{2.14}$$

Similarly we have

$$\begin{aligned} E[I_{3.2}] &\leq C\int_t^T ds\int_s^{s^+} du E[|Y_{s^+}^n||\dot{B}_u^n||\dot{B}_s^n|]ds \\ &\leq C\int_t^T ds\int_s^{s^+} du (E[(Y_{s^+}^n)^2])^{\frac{1}{2}}(E[|\dot{B}_u^n|^2|\dot{B}_s^n|^2])^{\frac{1}{2}}ds \\ &\leq C\int_t^T ds\int_s^{s^+} du (E[(Y_{s^+}^n)^2])^{\frac{1}{2}}2^n \\ &\leq C\int_t^T E[(Y_{s^+}^n)^2]ds + C. \end{aligned} \tag{2.15}$$

By virtue of the independence of $Y^n_{s^+}$ and $\dot{B}^n_s$, we have

$$\begin{aligned}
E[I_{3.3}] &\leq CE[\int_t^T |Y^n_{s^+}||\dot{B}^n_s|| \int_s^{s^+} Z^n_u dW_u|ds] \\
&\leq C\int_t^T (E[(Y^n_{s^+})^2|\dot{B}^n_s|^2])^{\frac{1}{2}}(E[|\int_s^{s^+} Z^n_u dW_u|^2])^{\frac{1}{2}}ds \\
&= C\int_t^T (E[(Y^n_{s^+})^2])^{\frac{1}{2}}(E[|\dot{B}^n_s|^2])^{\frac{1}{2}}(E[\int_s^{s^+} (Z^n_u)^2 du])^{\frac{1}{2}}ds \\
&= C\int_t^T (E[(Y^n_{s^+})^2])^{\frac{1}{2}}(2^n)^{\frac{1}{2}}(E[\int_s^{s^+} (Z^n_u)^2 du])^{\frac{1}{2}}ds \\
&\leq \frac{1}{4}\int_t^T 2^n E[\int_s^{s^+} (Z^n_u)^2 du]ds + C_2\int_t^T E[(Y^n_{s^+})^2]ds \\
&= \frac{1}{4}E[\int_t^T (Z^n_u)^2 du 2^n(\int_{u^-}^u ds)] + C_2\int_t^T E[(Y^n_{s^+})^2]ds \\
&\leq \frac{1}{4}E[\int_t^T (Z^n_u)^2 du] + C_2\int_t^T E[(Y^n_{s^+})^2]ds \qquad (2.16)
\end{aligned}$$

(2.14)–(2.16) imply that

$$E[I_3] \leq \frac{1}{4}E[\int_t^T (Z^n_u)^2 du] + C_2\int_t^T E[(Y^n_{s^+})^2]ds + C \qquad (2.17)$$

It follows from (2.5),(2.6),(2.12) and (2.17) that

$$\begin{aligned}
&E[(Y^n_t)^2] + \frac{1}{2}E[\int_t^T (Z^n_s)^2 ds] \\
\leq\ & E[(\xi)^2] + C\int_t^T E[(Y^n_{s^+})^2]ds + C \qquad (2.18)
\end{aligned}$$

Applying the Gronwall's inequality, we complete the proof of the Proposition.

The above result can be strengthened as

**Proposition 2.2** *For any $p \geq 1$, there exists a constant $C$ such that*

$$\sup_n\{E[\sup_{0\leq t\leq T} |Y^n_t|^p] + E[(\int_t^T (Z^n_s)^2 ds)^p]\} \leq C. \qquad (2.19)$$

**Theorem 2.3**

$$\lim_{n\to\infty}\sup_{0\leq t\leq T}\{E[(Y^n_t - Y_t)^2] + E[\int_t^T (Z^n_s - Z_s)^2 ds]\} = 0. \qquad (2.20)$$

**Proof**. By Ito's formula, we have

$$\begin{aligned}
& (Y_t^n - Y_t)^2 + \int_t^T (Z_s^n - Z_s)^2 ds \\
= \; & 2\int_t^T (Y_s^n - Y_s)(f(Y_s^n, Z_s^n) - f(Y_s, Z_s))ds \\
+ \; & 2\int_t^T (Y_s^n - Y_s)g(Y_s^n)dB_s^n + \int_t^T g^2(Y_s)ds \\
- \; & 2\int_t^T (Y_s^n - Y_s)g(Y_s)dB_s - 2\int_t^T (Y_s^n - Y_s)(Z_s^n - Z_s)dW_s \\
- \; & \int_t^T (Y_s^n - Y_s)gg'(Y_s)ds \\
:= \; & I_1^n + I_2^n + I_3^n + I_4^n + I_5^n + I_6^n. \qquad (2.21)
\end{aligned}$$

Crucially we need to bound the term $I_2^n$. We write

$$\begin{aligned}
I_2^n &= 2\int_t^T (Y_s^n - Y_s)g(Y_s^n)dB_s^n \\
&= 2\int_t^T [(Y_s^n - Y_s) - (Y_{s^+}^n - Y_{s^+})]g(Y_s^n)dB_s^n \\
&+ 2\int_t^T (Y_{s^+}^n - Y_{s^+})(g(Y_s^n) - g(Y_{s^+}^n))dB_s^n \\
&+ 2\int_t^T (Y_{s^+}^n - Y_{s^+})g(Y_{s^+}^n)dB_s^n \\
&:= A + B + C. \qquad (2.22)
\end{aligned}$$

As a stochastic integral, we have $E[C] = 0$. It is a long proof to establish the bounds for $E[A]$ and $E[B]$. We will split it into two lemmas for clarity.

**Lemma 2.4** *We have*

$$\begin{aligned}
E[A] \; \leq \; & C(\frac{1}{2^n})^{\frac{1}{2}-\delta} - 2E[\int_t^T g(Y_{s^+})g(Y_{s^+}^n)ds] \\
& + E[\int_t^T g^2(Y_{s^+}^n)ds]. \qquad (2.23)
\end{aligned}$$

**Proof**. By the equations satisfied by $Y^n$ and $Y$ we have

$$\begin{aligned} A &= 2\int_t^T (\int_s^{s^+} g(Y_u^n)dB_u^n)g(Y_s^n)dB_s^n \\ &- 2\int_t^T (\int_s^{s^+} g(Y_u)dB_u)g(Y_s^n)dB_s^n \\ &+ 2\int_t^T (\int_s^{s^+} [f(Y_u^n, Z_u^n) - f(Y_u, Z_u)]du)g(Y_s^n)dB_s^n \\ &- 2\int_t^T (\int_s^{s^+} (Z_u^n - Z_u)dW_u)g(Y_s^n)dB_s^n \\ &- \int_t^T (\int_s^{s^+} gg'(Y_u)du)g(Y_s^n)dB_s^n \\ &:= A_1 + A_2 + A_3 + A_4 + A_5. \end{aligned} \tag{2.24}$$

Clearly,

$$E[|A_5|] \le C\frac{1}{2^n}\int_t^T E[|\dot{B}_s^n|]ds \le C(\frac{1}{2^n})^{\frac{1}{2}}, \tag{2.25}$$

also

$$E[|A_3|] \le C(\frac{1}{2^n})^{\frac{1}{2}}. \tag{2.26}$$

By conditioning on $\mathcal{F}_s$, we find that

$$\begin{aligned} E[A_4] &= -2\int_t^T E\left[(\int_s^{s^+} (Z_u^n - Z_u)dW_u)g(Y_s^n)\dot{B}_s^n\right]ds \\ &= -2\int_t^T E\left[E[(\int_s^{s^+} (Z_u^n - Z_u)dW_u)|\mathcal{F}_s]g(Y_s^n)\dot{B}_s^n\right]ds \\ &= 0. \end{aligned} \tag{2.27}$$

To bound $A_1$, we write it as

$$\begin{aligned} A_1 &= 2\int_t^T (\int_s^{s^+} g(Y_u^n)dB_u^n)g(Y_s^n)dB_s^n \\ &= 2\int_t^T [\int_s^{s^+} (g(Y_u^n) - g(Y_{s^+}^n))dB_u^n]g(Y_s^n)dB_s^n \\ &+ 2\int_t^T g(Y_{s^+}^n)(B_s^n - B_{s^+}^n)(g(Y_s^n) - g(Y_{s^+}^n))dB_s^n \\ &+ 2\int_t^T g(Y_{s^+}^n)(B_s^n - B_{s^+}^n)g(Y_{s^+}^n)dB_s^n \\ &:= A_{11} + A_{12} + A_{13}. \end{aligned} \tag{2.28}$$

Splitting the interval $[t,T]$ into subintervals $[\frac{k}{2^n},\frac{k+1}{2^n}]$ we see that

$$\begin{aligned} A_{13} &= 2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} g^2(Y^n_{\frac{k+2}{2^n}})2^n(B_{\frac{k+2}{2^n}}-B_{\frac{k+3}{2^n}})(B_{\frac{k+2}{2^n}}-B_{\frac{k+1}{2^n}})\overrightarrow{ds} \\ &+ 2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} g^2(Y^n_{\frac{k+2}{2^n}})(2^n)^2(s-\frac{k+1}{2^n})(B_{\frac{k+2}{2^n}}-B_{\frac{k+1}{2^n}})^2\overrightarrow{ds} \\ &:= A_{13,1}+A_{13,2}. \end{aligned} \tag{2.29}$$

Conditioning on $\mathcal{F}_{\frac{k+2}{2^n}}$ we have

$$\begin{aligned} E[A_{13,1}] &= -2\sum_k E[g^2(Y^n_{\frac{k+2}{2^n}})(B_{\frac{k+2}{2^n}}-B_{\frac{k+3}{2^n}})E[(B_{\frac{k+2}{2^n}}-B_{\frac{k+1}{2^n}})|\mathcal{F}_{\frac{k+2}{2^n}}]] \\ &= 0. \end{aligned} \tag{2.30}$$

Integrating with respect to $s$, we get that

$$\begin{aligned} & E[A_{13}] = E[A_{13,2}] \\ &= E[\sum_k g^2(Y^n_{\frac{k+2}{2^n}})(B_{\frac{k+2}{2^n}}-B_{\frac{k+1}{2^n}})^2] \\ &= E[\sum_k g^2(Y^n_{\frac{k+2}{2^n}})\{(B_{\frac{k+2}{2^n}}-B_{\frac{k+1}{2^n}})^2-\frac{1}{2^n}\}] \\ &\qquad +E[\int_t^T g^2(Y^n_{s^+})ds] \\ &= E[\int_t^T g^2(Y^n_{s^+})ds], \end{aligned} \tag{2.31}$$

where the fact that the sequence $\{(B_{\frac{k+2}{2^n}}-B_{\frac{k+1}{2^n}})^2-\frac{1}{2^n},k\geq 0\}$ is a martingale has been used. For the term $A_{11}$ in (2.28), we have

$$\begin{aligned} A_{11} &= 2\int_0^1 d\lambda\int_t^T\int_s^{s^+} g'(Y^n_{s^+}+\lambda(Y^n_u-Y^n_{s^+}))(Y^n_u-Y^n_{s^+})dB^n_u g(Y^n_s)dB^n_s \\ &= 2\int_0^1 d\lambda\int_t^T\int_s^{s^+} g'(Y^n_{s^+}+\lambda(Y^n_u-Y^n_{s^+}))[\int_u^{s^+} f(Y^n_v,Z^n_v)dv]dB^n_u g(Y^n_s)dB^n_s \\ &+ 2\int_0^1 d\lambda\int_t^T\int_s^{s^+} g'(Y^n_{s^+}+\lambda(Y^n_u-Y^n_{s^+}))[\int_u^{s^+} g(Y^n_v)dB^n_v]dB^n_u g(Y^n_s)dB^n_s \\ &- 2\int_0^1 d\lambda\int_t^T\int_s^{s^+} g'(Y^n_{s^+}+\lambda(Y^n_u-Y^n_{s^+}))[\int_u^{s^+} Z^n_v dW_v]dB^n_u g(Y^n_s)dB^n_s \\ &:= A_{11,1}+A_{11,2}+A_{11,3}. \end{aligned} \tag{2.32}$$

The first two terms on the right can be bounded as follows.

$$\begin{aligned} E[A_{11,1}] &\leq C\frac{1}{2^n}\int_t^T ds\int_s^{s^+} E[|\dot{B}^n_u||\dot{B}^n_s|]du \\ &\leq C\frac{1}{2^n}(2^n)^{\frac{1}{2}}(2^n)^{\frac{1}{2}}\frac{1}{2^n}\leq C\frac{1}{2^n}. \end{aligned} \tag{2.33}$$

$$
\begin{aligned}
E[A_{11,2}] &\leq C\int_t^T ds\int_s^{s^+} du\int_u^{s^+} dvE[|\dot{B}^n_u||\dot{B}^n_v||\dot{B}^n_s|] \\
&\leq C(2^n)^{\frac{3}{2}}(\frac{1}{2^n})^2 \leq C(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.34)
\end{aligned}
$$

The last term $A_{11,3}$ can be estimated as follows.

$$
\begin{aligned}
E[A_{11,3}] &\leq CE[\int_t^T ds\int_s^{s^+} |\dot{B}^n_u||\dot{B}^n_s||\int_u^{s^+} Z^n_v dW_v|du] \\
&\leq C\int_t^T ds\int_s^{s^+} du(E[|\dot{B}^n_u|^2|\dot{B}^n_s|^2])^{\frac{1}{2}}(E[|\int_u^{s^+} Z^n_v dW_v|^2])^{\frac{1}{2}} \\
&\leq C2^n\int_t^T ds\int_s^{s^+} du(E[\int_u^{s^+}(Z^n_v)^2dv])^{\frac{1}{2}} \\
&\leq C\int_t^T ds(E[\int_s^{s^+}(Z^n_v)^2dv])^{\frac{1}{2}} \\
&\leq C\left(\int_t^T dsE[\int_s^{s^+}(Z^n_v)^2dv]\right)^{\frac{1}{2}} \leq C\left(E[\int_t^T (Z^n_v)^2dv\int_{v^-}^v ds]\right)^{\frac{1}{2}} \\
&\leq C(\sup_n(E[\int_t^T (Z^n_v)^2dv)^{\frac{1}{2}})(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.35)
\end{aligned}
$$

Putting together (2.33)–(2.35) together we get

$$
E[A_{11}] \leq C(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.36)
$$

Similarly the term $A_{12}$ can be decomposed as

$$
\begin{aligned}
A_{12} &= 2\int_0^1 d\lambda\int_t^T g(Y^n_{s^+})\int_s^{s^+} dB^n_u g'(Y^n_{s^+}+\lambda(Y^n_s-Y^n_{s^+}))(Y^n_s-Y^n_{s^+})dB^n_s \\
&= 2\int_0^1 d\lambda\int_t^T g(Y^n_{s^+})\int_s^{s^+} dB^n_u g'(Y^n_{s^+}+\lambda(Y^n_s-Y^n_{s^+}))[\int_s^{s^+} f(Y^n_v,Z^n_v)dv]dB^n_s \\
&+ 2\int_0^1 d\lambda\int_t^T g(Y^n_{s^+})\int_s^{s^+} dB^n_u g'(Y^n_{s^+}+\lambda(Y^n_s-Y^n_{s^+}))[\int_s^{s^+} g(Y^n_v)dB^n_v]dB^n_s \\
&- 2\int_0^1 d\lambda\int_t^T g(Y^n_{s^+})\int_s^{s^+} dB^n_u g'(Y^n_{s^+}+\lambda(Y^n_s-Y^n_{s^+}))[\int_s^{s^+} Z^n_v dW_v]dB^n_s \\
&:= A_{12,1}+A_{12,2}+A_{12,3}. \qquad (2.37)
\end{aligned}
$$

Using the similar arguments as for (2.33) and (2.34) we can show that

$$
E[A_{12,j}] \leq C\frac{1}{2^n}, j=1,2,3. \qquad (2.38)
$$

Hence,

$$
E[A_{12}] \leq C(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.39)
$$

Combining (2.31),(2.36) and (2.39) we get

$$E[A_1] \leq C(\frac{1}{2^n})^{\frac{1}{2}} + E[\int_t^T g^2(Y^n_{s^+})ds]. \tag{2.40}$$

Now we turn to $A_2$ which can be written as

$$\begin{aligned} A_2 &= -2\int_t^T (\int_s^{s^+} g(Y_u)dB_u)g(Y^n_s)dB^n_s \\ &= -2\int_t^T [\int_s^{s^+} (g(Y_u) - g(Y_{s^+}))dB_u]g(Y^n_s)dB^n_s \\ &- 2\int_t^T g(Y_{s^+})(B_s - B_{s^+})(g(Y^n_s) - g(Y^n_{s^+}))dB^n_s \\ &- 2\int_t^T g(Y_{s^+})(B_s - B_{s^+})g(Y^n_{s^+})dB^n_s \\ &:= A_{21} + A_{22} + A_{23}. \end{aligned} \tag{2.41}$$

Splitting the interval $[t, T]$ into subintervals $[\frac{k}{2^n}, \frac{k+1}{2^n}]$,

$$\begin{aligned} A_{23} &= -2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} g(Y_{\frac{k+2}{2^n}})g(Y^n_{\frac{k+2}{2^n}})(B_s - B_{\frac{k+2}{2^n}})2^n(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})\overrightarrow{ds} \\ &= -2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} g(Y_{\frac{k+2}{2^n}})g(Y^n_{\frac{k+2}{2^n}})(B_s - B_{\frac{k+1}{2^n}})2^n(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})\overrightarrow{ds} \\ &- 2\sum_k g(Y_{\frac{k+2}{2^n}})g(Y^n_{\frac{k+2}{2^n}})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})^2 \\ &:= A_{23,1} + A_{23,2}. \end{aligned} \tag{2.42}$$

Using the independence of the increments of $B$ and conditioning on $\mathcal{F}_{\frac{k+2}{2^n}}$ it follows that

$$\begin{aligned} & E[A_{23,1}] \\ =\ & -2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} 2^n\overrightarrow{ds}E[g(Y_{\frac{k+2}{2^n}})g(Y^n_{\frac{k+2}{2^n}})E[(B_s - B_{\frac{k+1}{2^n}})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})|\mathcal{F}_{\frac{k+2}{2^n}}]] \\ =\ & -2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} 2^n\overrightarrow{ds}E[g(Y_{\frac{k+2}{2^n}})g(Y^n_{\frac{k+2}{2^n}})E[(B_s - B_{\frac{k+1}{2^n}})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})]] \\ =\ & 0. \end{aligned} \tag{2.43}$$

Now,

$$\begin{aligned} A_{23,2} &= -2\sum_k g(Y_{\frac{k+2}{2^n}})g(Y^n_{\frac{k+2}{2^n}})\{(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})^2 - \frac{1}{2^n}\} \\ &- 2\int_t^T g(Y_{s^+})g(Y^n_{s^+})ds \end{aligned} \tag{2.44}$$

By conditioning on $\mathcal{F}_{\frac{k+2}{2^n}}$ we see that the expectation of the first term of the above equation vanishes. Hence,

$$E[A_{23}] = E[A_{23,2}] = -2E[\int_t^T g(Y_{s^+})g(Y^n_{s^+})ds]. \tag{2.45}$$

For the term $A_{21}$ we have

$$\begin{aligned}
A_{21} &= -2\int_0^1 d\lambda \int_t^T \int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+}))(Y_u - Y_{s^+})dB_u g(Y^n_s)dB^n_s \\
&= -2\int_0^1 d\lambda \int_t^T \int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+}))[\int_u^{s^+} f(Y_v, Z_v)dv]dB_u g(Y^n_s)dB^n_s \\
&- 2\int_0^1 d\lambda \int_t^T \int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+}))[\int_u^{s^+} g(Y_v)dB_v]dB_u g(Y^n_s)dB^n_s \\
&+ 2\int_0^1 d\lambda \int_t^T \int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+}))[\int_u^{s^+} Z_v dW_v]dB_u g(Y^n_s)dB^n_s \\
&:= A_{21,1} + A_{21,2} + A_{21,3}.
\end{aligned} \tag{2.46}$$

We will estimate each of the terms on the right. First we have

$$\begin{aligned}
E[A_{21,1}] &\le C\int_0^1 d\lambda \int_t^T E\left[|\int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+}))[\int_u^{s^+} f(Y_v, Z_v)dv]dB_u||\dot{B}^n_s|\right] ds \\
&\le C\int_0^1 d\lambda \int_t^T (E[|\int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+})) \\
&\qquad \times[\int_u^{s^+} f(Y_v, Z_v)dv]dB_u|^2])^{\frac{1}{2}}(E[|\dot{B}^n_s|^2])^{\frac{1}{2}}ds \\
&\le C(2^n)^{\frac{1}{2}}\int_0^1 d\lambda \int_t^T (E[\int_s^{s^+} g'(Y_{s^+} + \lambda(Y_u - Y_{s^+}))^2(\int_u^{s^+} f(Y_v, Z_v)dv)^2 du])^{\frac{1}{2}}ds \\
&\le C(2^n)^{\frac{1}{2}}\int_t^T [(s^+ - s)(s^+ - s)^2]^{\frac{1}{2}}ds \\
&\le C(\frac{1}{2^n}).
\end{aligned} \tag{2.47}$$

Similarly,

$$
\begin{aligned}
& E[A_{21,2}] \\
\leq\ & C\int_0^1 d\lambda \int_t^T E\left[ |\int_s^{s^+} g'(Y_{s^+}+\lambda(Y_u-Y_{s^+}))[\int_u^{s^+} g(Y_v)dB_v]dB_u||\dot{B}_s^n| \right] ds \\
\leq\ & C\int_0^1 d\lambda \int_t^T (E[|\int_s^{s^+} g'(Y_{s^+}+\lambda(Y_u-Y_{s^+}))[\int_u^{s^+} g(Y_v)dB_v]dB_u|^2])^{\frac{1}{2}}(E[|\dot{B}_s^n|^2])^{\frac{1}{2}}ds \\
\leq\ & C(2^n)^{\frac{1}{2}}\int_0^1 d\lambda \int_t^T (E[\int_s^{s^+} g'(Y_{s^+}+\lambda(Y_u-Y_{s^+}))^2(\int_u^{s^+} g(Y_v)dB_v)^2du])^{\frac{1}{2}}ds \\
\leq\ & C(2^n)^{\frac{1}{2}}\int_t^T (\int_s^{s^+} E[(\int_u^{s^+} g(Y_v)dB_v)^2]du)^{\frac{1}{2}}ds \\
\leq\ & C(2^n)^{\frac{1}{2}}\int_t^T (\int_s^{s^+} du)^{\frac{1}{2}}(E[\int_s^{s^+} g^2(Y_v)dv)])^{\frac{1}{2}}ds \\
\leq\ & C(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.48)
\end{aligned}
$$

By Hölder inequality and Ito isometry, we have

$$
\begin{aligned}
& E[A_{21,3}] \\
\leq\ & C\int_0^1 d\lambda \int_t^T E\left[ |\int_s^{s^+} g'(Y_{s^+}+\lambda(Y_u-Y_{s^+}))[\int_u^{s^+} Z_v dW_v]dB_u||\dot{B}_s^n| \right] ds \\
\leq\ & C\int_0^1 d\lambda \int_t^T (E[|\int_s^{s^+} g'(Y_{s^+}+\lambda(Y_u-Y_{s^+}))[\int_u^{s^+} Z_v dW_v]dB_u|^2])^{\frac{1}{2}}(E[|\dot{B}_s^n|^2])^{\frac{1}{2}}ds \\
\leq\ & C(2^n)^{\frac{1}{2}}\int_0^1 d\lambda \int_t^T (E[\int_s^{s^+} g'(Y_{s^+}+\lambda(Y_u-Y_{s^+}))^2(\int_u^{s^+} Z_v dW_v)^2du])^{\frac{1}{2}}ds \\
\leq\ & C(2^n)^{\frac{1}{2}}\int_t^T (\int_s^{s^+} E[(\int_u^{s^+} Z_v dW_v)^2]du)^{\frac{1}{2}}ds \\
\leq\ & C(2^n)^{\frac{1}{2}}\int_t^T (\int_s^{s^+} du)^{\frac{1}{2}}(E[\int_s^{s^+} Z_v^2 dv])^{\frac{1}{2}}ds \\
\leq\ & C(\int_t^T (E[\int_s^{s^+} Z_v^2 dv]ds)^{\frac{1}{2}} \\
\leq\ & C(E[\int_t^T Z_v^2 dv \int_{v^-}^v ds])^{\frac{1}{2}} \\
\leq\ & C(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.49)
\end{aligned}
$$

It follows from (2.47), (2.48) and (2.49)that

$$E[A_{21}] \leq C(\frac{1}{2^n})^{\frac{1}{2}}. \qquad (2.50)$$

Let us turn to the term $A_{22}$. We have

$$\begin{aligned} A_{22} &\leq C\int_t^T |B_s - B_{s^+}||Y_s^n - Y_{s^+}^n||\dot{B}_s^n|ds \\ &\leq C\int_t^T |B_s - B_{s^+}||\int_s^{s^+} f(Y_u^n, Z_u^n)du||\dot{B}_s^n|ds \\ &+ C\int_t^T |B_s - B_{s^+}||\int_s^{s^+} g(Y_u^n)dB_u^n||\dot{B}_s^n|ds \\ &+ C\int_t^T |B_s - B_{s^+}||\int_s^{s^+} Z_u^n dW_u||\dot{B}_s^n|ds \\ &:= A_{22,1} + A_{22,2} + A_{22,3}. \end{aligned} \tag{2.51}$$

Now,

$$\begin{aligned} E[A_{22,1}] &\leq C\frac{1}{2^n}\int_t^T (E[|B_s - B_{s^+}|^2])^{\frac{1}{2}}(E[|\dot{B}_s^n|^2])^{\frac{1}{2}}ds \\ &\leq C\frac{1}{2^n}, \end{aligned} \tag{2.52}$$

and

$$\begin{aligned} E[A_{22,2}] &\leq C\frac{1}{2^n}\int_t^T (E[|B_s - B_{s^+}|^2])^{\frac{1}{2}}(E[\sup_u |\dot{B}_u^n|^4])^{\frac{1}{2}}ds \\ &\leq C(\frac{1}{2^n})^{\frac{1}{2}}(2^n)^2\int_t^T (E[\sup_{|r-v|\leq\frac{1}{2^n}} |B_r - B_v|^4])^{\frac{1}{2}}ds \\ &\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \end{aligned} \tag{2.53}$$

By Hölder inequality,

$$\begin{aligned} E[A_{22,3}] &\leq C\int_t^T (E[|B_s - B_{s^+}|^4])^{\frac{1}{4}}(E[|\dot{B}_s^n|^4])^{\frac{1}{4}}(E[|\int_s^{s^+} Z_u^n dW_u|^2])^{\frac{1}{2}}ds \\ &\leq C(\int_t^T E[\int_s^{s^+} (Z_u^n)^2 du]ds)^{\frac{1}{2}} \\ &\leq C\frac{1}{2^n}. \end{aligned} \tag{2.54}$$

It follows from (2.51)–(2.54) that

$$E[A_{22}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.55}$$

Collecting (2.45),(2.50) and (2.51) we obtain

$$E[A_2] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta} - 2E[\int_t^T g(Y_{s^+})g(Y_{s^+}^n)ds]. \tag{2.56}$$

The proof is now completed by putting (2.25),(2.26),(2.27),(2.40) and (2.56) together.

**Lemma 2.5** *We have*

$$E[B] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta} + E[\int_t^T (Y^n_{s^+} - Y_{s^+}) g g'(Y^n_{s^+}) ds]. \tag{2.57}$$

**Proof**. Write

$$\begin{aligned}
B &= 2\int_t^T (Y^n_{s^+} - Y_{s^+})(g(Y^n_s) - g(Y^n_{s^+}))dB^n_s \\
&= 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+} + \lambda(Y^n_s - Y^n_{s^+}))(Y^n_s - Y^n_{s^+})dB^n_s \\
&= 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+} + \lambda(Y^n_s - Y^n_{s^+}))[\int_s^{s^+} f(Y^n_v, Z^n_v)dv]dB^n_s \\
&+ 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+} + \lambda(Y^n_s - Y^n_{s^+}))[\int_s^{s^+} g(Y^n_v)dB^n_v]dB^n_s \\
&- 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+} + \lambda(Y^n_s - Y^n_{s^+}))[\int_s^{s^+} Z^n_v dW_v]dB^n_s \\
&:= B_1 + B_2 + B_3.
\end{aligned} \tag{2.58}$$

By Proposition 2.2, we have

$$\begin{aligned}
E[B_1] &\leq C\frac{1}{2^n} E[\int_t^T |Y^n_{s^+} - Y_{s^+}||\dot{B}^n_s|ds] \\
&\leq C\frac{1}{2^n}\int_t^T (E[|Y^n_{s^+} - Y_{s^+}|^2])^{\frac{1}{2}} (E[|\dot{B}^n_s|^2])^{\frac{1}{2}} ds \\
&\leq C(\frac{1}{2^n})^{\frac{1}{2}}.
\end{aligned} \tag{2.59}$$

$B_2$ is further written as follows.

$$\begin{aligned}
B_2 &= 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+})[g'(Y^n_{s^+} + \lambda(Y^n_s - Y^n_{s^+})) - g'(Y^n_{s^+})][\int_s^{s^+} g(Y^n_u)dB^n_u]dB^n_s \\
&+ 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+})[\int_s^{s^+} (g(Y^n_u) - g(Y^n_{s^+}))dB^n_u]dB^n_s \\
&+ 2\int_0^1 d\lambda \int_t^T (Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+}) g(Y^n_{s^+})(B^n_s - B^n_{s^+})dB^n_s \\
&:= B_{21} + B_{22} + B_{23}.
\end{aligned} \tag{2.60}$$

By the Lipschitz continuity of $g'$, it follows that

$$\begin{aligned}
B_{21} &\leq C\int_t^T |Y^n_{s^+}-Y_{s^+}||Y^n_s-Y^n_{s^+}|\Big|\int_s^{s^+} g(Y^n_u)dB^n_u\Big||\dot{B}^n_s|ds\\
&\leq C\int_t^T |Y^n_{s^+}-Y_{s^+}|\Big|\int_s^{s^+} f(Y^n_v,Z^n_v)dv\Big|\Big|\int_s^{s^+} g(Y^n_u)dB^n_u\Big||\dot{B}^n_s|ds\\
&+ C\int_t^T |Y^n_{s^+}-Y_{s^+}|\Big|\int_s^{s^+} g(Y^n_u)dB^n_u|^2|\dot{B}^n_s|ds\\
&+ C\int_t^T |Y^n_{s^+}-Y_{s^+}|\Big|\int_s^{s^+} Z^n_v dW_v\Big|\Big|\int_s^{s^+} g(Y^n_u)dB^n_u\Big||\dot{B}^n_s|ds\\
&:= B_{21,1}+B_{21,2}+B_{21,3}. \qquad (2.61)
\end{aligned}$$

The following two inequalities will be used frequently in sequel.

$$\sup_u|\dot{B}^n_u|\leq 2^n \sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s| \qquad (2.62)$$

For any $\delta>0$ and $p\geq 1$, there exists a constant $C_{p,\delta}$ such that

$$E[\sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s|^p]\leq C_{p,\delta}(\frac{1}{2^n})^{\frac{p}{2}-\delta}. \qquad (2.63)$$

By Hölder's inequality and (2.62), (2.63), we have

$$\begin{aligned}
E[B_{21,1}] &\leq C\frac{1}{2^n}E[\int_t^T |Y^n_{s^+}-Y_{s^+}|\int_s^{s^+}|\dot{B}^n_u|du||\dot{B}^n_s|ds]\\
&\leq C\frac{1}{2^n}(2^n)^2E[\int_t^T |Y^n_{s^+}-Y_{s^+}|\Big|\int_s^{s^+}du\Big|\sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s|^2ds]\\
&\leq CE[\int_t^T |Y^n_{s^+}-Y_{s^+}|\sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s|^2ds]\\
&\leq C\int_t^T (E[|Y^n_{s^+}-Y_{s^+}|^2])^{\frac{1}{2}}(E[\sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s|^4])^{\frac{1}{2}}ds\\
&\leq C(\frac{1}{2^n})^{1-\delta}. \qquad (2.64)
\end{aligned}$$

Similarly, in view of (2.63), we have

$$\begin{aligned}
E[B_{21,2}] &\leq CE[\int_t^T |Y^n_{s^+}-Y_{s^+}|\Big|\int_s^{s^+}du\Big|^2\sup_s|\dot{B}^n_s|^3ds]\\
&\leq C(\frac{1}{2^n})^2(2^n)^3E[\int_t^T |Y^n_{s^+}-Y_{s^+}|\sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s|^3ds]\\
&\leq C2^n\int_t^T (E[|Y^n_{s^+}-Y_{s^+}|^2])^{\frac{1}{2}}(E[\sup_{|r-s|\leq\frac{1}{2^n}}|B_r-B_s|^6])^{\frac{1}{2}}ds\\
&\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}, \qquad (2.65)
\end{aligned}$$

and

$$\begin{aligned}
E[B_{21,3}] &\leq CE[\int_t^T |Y^n_{s^+} - Y_{s^+}||\int_s^{s^+} Z^n_v dW_v|(\int_s^{s^+} |\dot{B}^n_u|du)|\dot{B}^n_s|ds] \\
&\leq C\frac{1}{2^n}(2^n)^2 E[\int_t^T |Y^n_{s^+} - Y_{s^+}||\int_s^{s^+} Z^n_v dW_v| \sup_{|r-s|\leq\frac{1}{2^n}} |B_r - B_s|^2 ds] \\
&\leq C2^n \int_t^T (E[|Y^n_{s^+} - Y_{s^+}|^4])^{\frac{1}{4}} (E[|\int_s^{s^+} Z^n_v dW_v|^2])^{\frac{1}{2}} (E[\sup_{|r-s|\leq\frac{1}{2^n}} |B_r - B_s|^8])^{\frac{1}{4}} ds \\
&\leq C(\frac{1}{2^n})^{1-\delta} 2^n \int_t^T (E[|Y^n_{s^+} - Y_{s^+}|^4])^{\frac{1}{4}} (E[\int_s^{s^+} (Z^n_v)^2 dv])^{\frac{1}{2}} ds \\
&\leq C(\frac{1}{2^n})^{1-\delta} 2^n (\int_t^T (E[|Y^n_{s^+} - Y_{s^+}|^4])^{\frac{1}{2}} ds)^{\frac{1}{2}} (\int_t^T E[\int_s^{s^+} (Z^n_v)^2 dv] ds)^{\frac{1}{2}} \\
&\leq C(\frac{1}{2^n})^{1-\delta} 2^n (E[\int_t^T (Z^n_v)^2 dv \int_{v^-}^v ds])^{\frac{1}{2}} \\
&\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}, \qquad (2.66)
\end{aligned}$$

where the a priori estimate (2.19) has been used. (2.64)–(2.64)yields

$$E[B_{21}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \qquad (2.67)$$

By the Lipschitz continuity of $g$, we have

$$\begin{aligned}
B_{22} &\leq C\int_t^T |Y^n_{s^+} - Y_{s^+}|(\int_s^{s^+} |Y^n_u - Y^n_{s^+}|\dot{B}^n_u|du)|\dot{B}^n_s|ds \\
&\leq C\int_t^T |Y^n_{s^+} - Y_{s^+}|(\int_s^{s^+} |\int_u^{s^+} f(Y^n_v, Z^n_v)dv||\dot{B}^n_u|du)|\dot{B}^n_s|ds \\
&\leq C\int_t^T |Y^n_{s^+} - Y_{s^+}|(\int_s^{s^+} |\int_u^{s^+} g(Y^n_v)dB^n_v||\dot{B}^n_u|du)|\dot{B}^n_s|ds \\
&\leq C\int_t^T |Y^n_{s^+} - Y_{s^+}|(\int_s^{s^+} |\int_u^{s^+} Z^n_v dW_v||\dot{B}^n_u|du)|\dot{B}^n_s|ds \\
&:= B_{22,1} + B_{22,2} + B_{22,3}. \qquad (2.68)
\end{aligned}$$

By the similar arguments as above, we have

$$E[B_{22,j}] \leq C(\frac{1}{2^n})^{1-\delta}, j = 1, 2, 3. \qquad (2.69)$$

Hence,

$$E[B_{22}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \qquad (2.70)$$

For the term $B_{23}$, we split the interval $[t,T]$ into subintervals $[\frac{k}{2^n},\frac{k+1}{2^n}]$ to get

$$\begin{aligned}
& B_{23} \\
= \;& 2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} (Y^n_{\frac{k+2}{2^n}} - Y_{\frac{k+2}{2^n}}) gg'(Y^n_{\frac{k+2}{2^n}}) 2^n (B_{\frac{k+2}{2^n}} - B_{\frac{k+3}{2^n}})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}}) \overrightarrow{ds} \\
+ \;& 2\sum_k \int_{\frac{k}{2^n}}^{\frac{k+1}{2^n}} (Y^n_{\frac{k+2}{2^n}} - Y_{\frac{k+2}{2^n}}) gg'(Y^n_{\frac{k+2}{2^n}}) (2^n)^2 (s - \frac{k+1}{2^n})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})^2 \overrightarrow{ds} \\
= \;& -2\sum_k (Y^n_{\frac{k+2}{2^n}} - Y_{\frac{k+2}{2^n}}) gg'(Y^n_{\frac{k+2}{2^n}})(B_{\frac{k+2}{2^n}} - B_{\frac{k+3}{2^n}})(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}}) \\
+ \;& \sum_k (Y^n_{\frac{k+2}{2^n}} - Y_{\frac{k+2}{2^n}}) gg'(Y^n_{\frac{k+2}{2^n}}) \{(B_{\frac{k+2}{2^n}} - B_{\frac{k+1}{2^n}})^2 - \frac{1}{2^n}\} \\
+ \;& \int_t^T (Y^n_{s^+} - Y_{s^+}) gg'(Y^n_{s^+}) ds \qquad (2.71)
\end{aligned}$$

Conditioning on $\mathcal{F}_{\frac{k+2}{2^n}}$ it is easy to see that the expectation of the first two terms on the right vanishes. Hence,

$$E[B_{23}] = E[\int_t^T (Y^n_{s^+} - Y_{s^+}) gg'(Y^n_{s^+}) ds]. \qquad (2.72)$$

Collect the terms in (2.67),(2.70) and (2.72) to obtain

$$E[B_2] \le C(\frac{1}{2^n})^{\frac{1}{2}-\delta} + E[\int_t^T (Y^n_{s^+} - Y_{s^+}) gg'(Y^n_{s^+}) ds]. \qquad (2.73)$$

Now we turn to the term $B_3$ in (2.58). We further split it as

$$\begin{aligned}
B_3 \;=\;& -2\{\int_0^1 d\lambda \int_t^T [(Y^n_{s^+} - Y_{s^+}) g'(Y^n_{s^+} + \lambda(Y^n_s - Y^n_{s^+})) \\
& -(Y^n_s - Y_s) g'(Y^n_s)](\int_s^{s^+} Z^n_v dW_v) dB^n_s\} \\
& -2\int_t^T (Y^n_s - Y_s) g'(Y^n_s)(\int_s^{s^+} Z^n_v dW_v) dB^n_s \\
:=\;& B_{31} + B_{32}. \qquad (2.74)
\end{aligned}$$

First we notice that

$$\begin{aligned}
E[B_{32}] \;=\;& -2\int_t^T E[(Y^n_s - Y_s) g'(Y^n_s) \dot{B}^n_s E[(\int_s^{s^+} Z^n_v dW_v)|\mathcal{F}_s]] ds \\
=\;& 0. \qquad (2.75)
\end{aligned}$$

By the Lipschitz continuity of $g'$, we have

$$\begin{aligned} B_{31} &\leq C\int_t^T |Y^n_{s^+} - Y^n_s|| \int_s^{s^+} Z^n_v dW_v||\dot{B}^n_s|ds \\ &+ C\int_t^T |Y_{s^+} - Y_s|| \int_s^{s^+} Z^n_v dW_v||\dot{B}^n_s|ds \\ &+ C\int_t^T |Y^n_s - Y_s||Y^n_{s^+} - Y^n_s|| \int_s^{s^+} Z^n_v dW_v||\dot{B}^n_s|ds \\ &:= B_{31,1} + B_{31,2} + B_{31,3}. \end{aligned} \tag{2.76}$$

Furthermore,

$$\begin{aligned} B_{31,1} &\leq C\int_t^T |\int_s^{s^+} f(Y^n_v, Z^n_v)dv|| \int_s^{s^+} Z^n_v dW_v||\dot{B}^n_s|ds \\ &+ C\int_t^T |\int_s^{s^+} g(Y^n_v)dB^n_v|| \int_s^{s^+} Z^n_v dW_v||\dot{B}^n_s|ds \\ &+ C\int_t^T |\int_s^{s^+} Z^n_v dW_v|^2|\dot{B}^n_s|ds \\ &:= B_{31,11} + B_{31,12} + B_{31,13}. \end{aligned} \tag{2.77}$$

Now, interchanging the order of integration, we have

$$\begin{aligned} E[B_{31,11}] &\leq C\frac{1}{2^n}\int_t^T (E[|\int_s^{s^+} (Z^n_v)^2 dv|^2])^{\frac{1}{2}} (E[|\dot{B}^n_s|^2])^{\frac{1}{2}} ds \\ &\leq C\frac{1}{2^n}. \end{aligned} \tag{2.78}$$

In view of (2.62), (2.63), we have

$$\begin{aligned} E[B_{31,12}] &\leq C\frac{1}{2^n}\int_t^T E[\sup_s |\dot{B}^n_s|^2| \int_s^{s^+} Z^n_v dW_v|]ds \\ &\leq C\frac{1}{2^n}(2^n)^2 \int_t^T (E[\sup_{|r-v|\leq \frac{1}{2^n}} |B_r - B_v|^4])^{\frac{1}{2}} (E[|\int_s^{s^+} Z^n_v dW_v|^2])^{\frac{1}{2}} ds \\ &\leq C\frac{1}{2^n}(2^n)^2 (\frac{1}{2^n})^{1-\delta} (\int_t^T E[\int_s^{s^+} (Z^n_v)^2 dv]ds)^{\frac{1}{2}} \\ &\leq C\frac{1}{2^n}(2^n)^2 (\frac{1}{2^n})^{1-\delta} (\frac{1}{2^n})^{\frac{1}{2}} (\int_t^T E[(Z^n_v)^2]dv)^{\frac{1}{2}} \\ &\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \end{aligned} \tag{2.79}$$

Using the fact that $|\dot{B}^n_s|$ is $\mathcal{F}_s$-measurable we have

$$\begin{aligned}
E[B_{31,13}] &\leq CE[\int_t^T (\int_s^{s^+} Z^n_v dW_v)^2 |\dot{B}^n_s| ds] \\
&= CE[\int_t^T (\int_s^{s^+} (|\dot{B}^n_s|)^{\frac{1}{2}} Z^n_v dW_v)^2 ds] \\
&= CE[\int_t^T \int_s^{s^+} (|\dot{B}^n_s|)(Z^n_v)^2 dv ds] \\
&\leq C(2^n) E[(\sup_{|r-v|\leq \frac{1}{2^n}} |B_r - B_v|) \int_t^T \int_s^{s^+} ((Z^n_v)^2 dv ds] \\
&\leq C(2^n)(E[(\sup_{|r-v|\leq \frac{1}{2^n}} |B_r - B_v|^2])^{\frac{1}{2}} (E[(\int_t^T \int_s^{s^+} (Z^n_v)^2 dv ds)^2])^{\frac{1}{2}} \\
&\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}, \qquad (2.80)
\end{aligned}$$

where (2.62),(2.63) again were used. (2.78)- (2.80) implies that

$$E[B_{31,1}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \qquad (2.81)$$

For the term $B_{31,2}$ we have

$$\begin{aligned}
B_{31,2} &\leq C \int_t^T |\int_s^{s^+} f(Y_v, Z_v) dv| |\int_s^{s^+} Z^n_v dW_v| |\dot{B}^n_s| ds \\
&+ C \int_t^T |\int_s^{s^+} g(Y_v) dB_v| |\int_s^{s^+} Z^n_v dW_v| |\dot{B}^n_s| ds \\
&+ C \int_t^T |\int_s^{s^+} Z_v dW_v| |\int_s^{s^+} Z^n_v dW_v| |\dot{B}^n_s| ds \\
&:= B_{31,21} + B_{31,22} + B_{31,23}. \qquad (2.82)
\end{aligned}$$

By a similar argument as for (2.78), we have

$$E[B_{31,21}] \leq C(\frac{1}{2^n}). \qquad (2.83)$$

As for $B_{31,22}$, we have

$$\begin{aligned}
E[B_{31,22}] &\leq C \int_t^T (E[|\int_s^{s^+} Z^n_u (|\dot{B}^n_s|)^{\frac{1}{2}} dW_u|^2])^{\frac{1}{2}} (E[|\int_s^{s^+} g(Y_u) dB_u|^2])^{\frac{1}{2}} ds \\
&\leq C(\frac{1}{2^n})^{\frac{1}{2}} \int_t^T (E[\int_s^{s^+} (Z^n_u)^2 |\dot{B}^n_s| du])^{\frac{1}{2}} ds \\
&\qquad (2.84)
\end{aligned}$$

From here following the same arguments as for (2.80) we obtain

$$E[B_{31,22}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-delta}. \qquad (2.85)$$

The term $B_{31,23}$ is bounded as

$$\begin{aligned} B_{31,23} &\leq C\int_t^T (\int_s^{s^+} Z_v^n dW_v)^2 |\dot{B}_s^n| ds \\ &+ C\int_t^T (\int_s^{s^+} Z_v dW_v)^2 |\dot{B}_s^n| ds, \end{aligned} \tag{2.86}$$

which, together with the same arguments as for (2.103), yields

$$E[B_{31,23}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.87}$$

(2.83)- (2.87) gives that

$$E[B_{31,2}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.88}$$

Finally we need to find an upper bound for $E[B_{31,3}]$. Notice that

$$\begin{aligned} B_{31,3} &\leq C\int_t^T |Y_s^n - Y_s|| \int_s^{s^+} f(Y_v^n, Z_v^n) dv|| \int_s^{s^+} Z_v^n dW_v||\dot{B}_s^n| ds \\ &+ C\int_t^T |Y_s^n - Y_s|| \int_s^{s^+} g(Y_v^n) dB_v^n|| \int_s^{s^+} Z_v^n dW_v||\dot{B}_s^n| ds \\ &+ C\int_t^T |Y_s^n - Y_s|| \int_s^{s^+} Z_v^n dW_v|^2 |\dot{B}_s^n| ds \\ &:= B_{31,31} + B_{31,32} + B_{31,33}. \end{aligned} \tag{2.89}$$

We have

$$\begin{aligned} E[B_{31,31}] &\leq C\frac{1}{2^n}\int_t^T (E[|Y_s^n - Y_s|^2])^{\frac{1}{2}} (E[|\int_s^{s^+} (|\dot{B}_s^n|)^{\frac{1}{2}} Z_v^n dW_v|^2])^{\frac{1}{2}} ds \\ &\leq C\frac{1}{2^n}(\int_t^T E[|Y_s^n - Y_s|^2] ds)^{\frac{1}{2}} (\int_t^T E[|\int_s^{s^+} |\dot{B}_s^n| (Z_v^n)^2 dv] ds)^{\frac{1}{2}} \\ &\leq C\frac{1}{2^n}(2^n)^{\frac{1}{2}} (E[(\sup_{|r-v|\leq \frac{1}{2^n}} |B_r - B_v|) \int_t^T \int_s^{s^+} (Z_v^n)^2 dv] ds)^{\frac{1}{2}} \\ &\leq C(\frac{1}{2^n})^{\frac{1}{2}} (E[\sup_{|r-v|\leq \frac{1}{2^n}} |B_r - B_v|^2])^{\frac{1}{4}} (E[(\int_t^T \int_s^{s^+} (Z_v^n)^2 dv ds)^2])^{\frac{1}{4}} \\ &\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta} (E[(\int_t^T (Z_v^n)^2 dv)^2])^{\frac{1}{4}} \\ &\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}, \end{aligned} \tag{2.90}$$

where the a priori bounds (2.4), (2.19) have been used. Noticing that $|Y_s^n - Y_s|, |\dot{B}_s^n|ds$ are $\mathcal{F}_s$ measurable we have

$$\begin{aligned}
E[B_{31,33}] &= C\int_t^T E[|Y_s^n - Y_s||\int_s^{s^+} Z_v^n dW_v|^2|\dot{B}_s^n|]ds \\
&= C\int_t^T E[|Y_s^n - Y_s|(\int_s^{s^+}(Z_v^n)^2dv)|\dot{B}_s^n|]ds \\
&\leq C2^n E[(\sup_{0\leq s\leq T}|Y_s^n - Y_s|)(\sup_{|r-v|\leq\frac{1}{2^n}}|B_r - B_v|^2])\int_t^T(\int_s^{s^+}(Z_v^n)^2dv)ds] \\
&\leq C2^n(E[\sup_{0\leq s\leq T}|Y_s^n - Y_s|^4])^{\frac{1}{4}}(E[\sup_{|r-v|\leq\frac{1}{2^n}}|B_r - B_v|^4])^{\frac{1}{4}} \\
&\qquad \times(E[(\int_t^T(\int_s^{s^+}(Z_v^n)^2dv)ds)^2])^{\frac{1}{2}} \\
&\leq C2^n(\frac{1}{2^n})^{\frac{1}{2}-\delta}(E[(\int_t^T(Z_v^n)^2(\int_{v^-}^v ds)dv)^2])^{\frac{1}{2}} \\
&\leq C2^n(\frac{1}{2^n})^{\frac{1}{2}-\delta}\frac{1}{2^n}(E[(\int_t^T(Z_v^n)^2dv)^2])^{\frac{1}{2}} \\
&\leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \qquad (2.91)
\end{aligned}$$

As for the term $B_{31.32}$ we have

$$\begin{aligned}
E[B_{31,32}] &\leq C\int_t^T E[|Y_s^n - Y_s||\int_s^{s^+} Z_v^n dW_v|^2|\dot{B}_s^n|]ds \\
&+ C\int_t^T E[|Y_s^n - Y_s||\int_s^{s^+} g(Y_v^n)dB_v^n|^2|\dot{B}_s^n|]ds. \qquad (2.92)
\end{aligned}$$

Furthermore, we have

$$\begin{aligned}
&\int_t^T E[|Y_s^n - Y_s||\int_s^{s^+} g(Y_v^n)dB_v^n|^2|\dot{B}_s^n|]ds \\
\leq\ & C(2^n)^3 E[(\sup_{|r-v|\leq\frac{1}{2^n}}|B_r - B_v|^3)\int_t^T|Y_s^n - Y_s|(\int_s^{s^+}|g(Y_v^n)|dv)^2ds] \\
\leq\ & C(2^n)(E[(\sup_{|r-v|\leq\frac{1}{2^n}}|B_r - B_v|^6])^{\frac{1}{2}}(E[(\int_t^T|Y_s^n - Y_s|ds)^2])^{\frac{1}{2}} \\
\leq\ & C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \qquad (2.93)
\end{aligned}$$

It follows now from (2.91), (2.92) and (2.93) that

$$E[B_{31,32}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \qquad (2.94)$$

(2.90)- (2.94) yields that

$$E[B_{31,3}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.95}$$

It follows from (2.81), (2.88) and (2.95) that

$$E[B_{31}] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.96}$$

This together with (2.75) yields

$$E[B_3] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.97}$$

The lemma now follows from (2.59), (2.73) and (2.97).

**Proof of Theorem 2.3**(continued).

We are ready to complete the proof of Theorem 2.3. Taking expectation in (2.21) we obtain

$$\begin{aligned}
& E[(Y_t^n - Y_t)^2] + E\left[\int_t^T (Z_s^n - Z_s)^2 ds\right] \\
= \;& 2E\left[\int_t^T (Y_s^n - Y_s)(f(Y_s^n, Z_s^n) - f(Y_s, Z_s))ds\right] \\
+ \;& 2E\left[\int_t^T (Y_s^n - Y_s)g(Y_s^n)dB_s^n\right] + E\left[\int_t^T g^2(Y_s)ds\right] \\
- \;& E\left[\int_t^T (Y_s^n - Y_s)gg'(Y_s)ds\right]
\end{aligned} \tag{2.98}$$

Taking into account of the estimates in Lemma 2.4 and Lemma 2.5 we deduce from (2.98) that

$$\begin{aligned}
& E[(Y_t^n - Y_t)^2] + E\left[\int_t^T (Z_s^n - Z_s)^2 ds\right] \\
\leq \;& C(\frac{1}{2^n})^{\frac{1}{2}-\delta} + CE\left[\int_t^T |Y_s^n - Y_s|^2 ds\right] \\
+ \;& \frac{1}{2}E\left[\int_t^T |Z_s^n - Z_s|^2 ds\right] + E\left[\int_t^T g^2(Y_s)ds\right] \\
- \;& E\left[\int_t^T (Y_s^n - Y_s)gg'(Y_s)ds\right] \\
+ \;& E[\int_t^T (Y_{s^+}^n - Y_{s^+})gg'(Y_{s^+}^n)ds] \\
& -2E[\int_t^T g(Y_{s^+})g(Y_{s^+}^n)ds] + E[\int_t^T g^2(Y_{s^+}^n)ds].
\end{aligned} \tag{2.99}$$

To proceed with the proof, we claim that there is a constant $C$ such that

$$E[\int_0^T |Y_{s^+}^n - Y_s^n|^2 ds] \leq C(\frac{1}{2^n})^{1-\delta}. \tag{2.100}$$

$$E[\int_0^T |Y_{s^+} - Y_s|^2 ds] \leq C(\frac{1}{2^n})^{1-\delta}. \tag{2.101}$$

Let us prove (2.100). The proof of (2.101) is similar. Indeed, we have

$$\begin{aligned}
& E[\int_0^T |Y_{s^+}^n - Y_s^n|^2 ds] \\
\leq\ & CE[\int_0^T |\int_s^{s^+} f(Y_u^n, Z_u^n) du|^2 ds] \\
+\ & CE[\int_0^T |\int_s^{s^+} g(Y_u^n) dB_u^n|^2 ds] \\
+\ & CE[\int_0^T |\int_s^{s^+} Z_u^n dW_u|^2 ds] \\
\leq\ & C\frac{1}{2^n} + CE[\sup_{|r-v|\leq \frac{1}{2^n}} |B_r - B_v|^2] \\
+\ & CE[\int_0^T \int_s^{s^+} (Z_u^n)^2 du ds] \\
\leq\ & C(\frac{1}{2^n})^{1-\delta} + \sup_n (E[\int_0^T (Z_u^n)^2 du]) \frac{1}{2^n} \\
\leq\ & C(\frac{1}{2^n})^{1-\delta}.
\end{aligned} \tag{2.102}$$

By Hölder inequality it follows immediately from (2.100) and (2.101) that

$$E[\int_0^T |Y_{s^+}^n - Y_s^n| ds] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.103}$$

$$E[\int_0^T |Y_{s^+} - Y_s| ds] \leq C(\frac{1}{2^n})^{\frac{1}{2}-\delta}. \tag{2.104}$$

Because of (2.100) and (2.101), we now replace $s^+$ by $s$ on the right side of (2.99) to obtain

$$\begin{aligned} & E[(Y_t^n - Y_t)^2] + E\left[\int_t^T (Z_s^n - Z_s)^2 ds\right] \\ \leq\ & C(\frac{1}{2^n})^{\frac{1}{2}-\delta} + CE\left[\int_t^T |Y_s^n - Y_s|^2 ds\right] \\ +\ & \frac{1}{2}E\left[\int_t^T |Z_s^n - Z_s|^2 ds\right] + E\left[\int_t^T g^2(Y_s) ds\right] \\ -\ & E\left[\int_t^T (Y_s^n - Y_s) g g'(Y_s) ds\right] \\ +\ & E[\int_t^T (Y_s^n - Y_s) g g'(Y_s^n) ds] \\ & -2E[\int_t^T g(Y_s) g(Y_s^n) ds] + E[\int_t^T g^2(Y_s^n) ds]. \end{aligned} \tag{2.105}$$

Remark that the constant $C$ in front of $(\frac{1}{2^n})^{\frac{1}{2}-\delta}$ is different from that in (2.99). Completing the square in (2.105) we get that

$$\begin{aligned} & E[(Y_t^n - Y_t)^2] + E\left[\int_t^T (Z_s^n - Z_s)^2 ds\right] \\ \leq\ & C(\frac{1}{2^n})^{\frac{1}{2}-\delta} + CE\left[\int_t^T |Y_s^n - Y_s|^2 ds\right] \\ +\ & \frac{1}{2}E\left[\int_t^T |Z_s^n - Z_s|^2 ds\right] + E\left[\int_t^T (g(Y_s) - g(Y_s^n))^2 ds\right] \\ +\ & E[\int_t^T (Y_s^n - Y_s)(g g'(Y_s^n) - g g'(Y_s)) ds]. \end{aligned} \tag{2.106}$$

By the Lipschitz continuity of $gg'$ and $g$, it follows from (2.106) that

$$\begin{aligned} & E[(Y_t^n - Y_t)^2] + \frac{1}{2}E\left[\int_t^T (Z_s^n - Z_s)^2 ds\right] \\ \leq\ & C(\frac{1}{2^n})^{\frac{1}{2}-\delta} + CE\left[\int_t^T |Y_s^n - Y_s|^2 ds\right]. \end{aligned} \tag{2.107}$$

Application of the Gronwall's inequality completes the proof of Theorem 2.3.

# References


[ES] L. C. Evans and D. W. Stroock: An approximation scheme for reflected stochastic differential equations, Stochastic Processes and Their applications 121 (2011) 1464-1491.

[IW] N. Ikeda and S. Watanable: Stochastic Differential Equations and Diffusion Processes. North-Holland, Amsterdam, 1981.

[Z] Tusheng Zhang: Wong-Zakai approximations to SDEs with reflection, to appear in Potential Analysis.

[PP1] E. Pardoux and S. Peng, Adapted solution of a backward stochastic differential equation.Systems Control Letter **14**(1990) 55-61.

[PP] E. Pardoux, S. Peng: Backward doubly SDEs and systems of quasilinear SPDEs, PTRF 98(1994) 209-227.

[WZ] E. Wong and M. Zakai: On the convegence of ordinary integrals to stochastic integrals, Ann. Math. Statist. 36(1965)1560-1564.

[WZ1] E. Wong and M. Zakai: On the relation between ordinary and stochastic differential equations, Internat. J. Engrg. Sci. 3 (1965) 213-229.